

## Results Of Evaluation Of Anti-Radical Properties Of *Inula Grandis Schrenk* (Asteraceae) Plant By DPPH Method

<sup>1</sup> Abdurashidova Gulnura Abdurashidovna

<sup>2</sup> Gulamov Ilhomjon Komiljonovich

<sup>3</sup> Haydarova Dilrabo Rahimjonovna

<sup>1</sup> Namangan State University, Master 2, Uzbekistan

<sup>2</sup> Tashkent University of Information Technologies, Associate Professor, PhD., Uzbekistan

<sup>3</sup> Namangan State University, Associate Professor, PhD., Uzbekistan

Received: 30<sup>th</sup> Nov 2025 | Received Revised Version: 18<sup>th</sup> Dec 2025 | Accepted: 02<sup>th</sup> Jan 2026 | Published: 25<sup>th</sup> Jan 2026

Volume 08 Issue 01 2026 | Crossref DOI: 10.37547/tajabe/Volume08Issue01-04

### Abstract

As a result of the conducted studies, it was found that alcoholic and aqueous extracts prepared from the *Inula grandis* plant have significant antiradical activity against DPPH free radicals. Spectrophotometric analyses based on the DPPH method showed a consistent increase in antiradical activity with increasing extract volume and incubation time. Based on the results obtained, the scientific and practical significance of the *Inula grandis* L. plant as a natural antioxidant source is demonstrated.

Keywords: *Inula grandis*, phytochemistry, flora, DPPH, antiradical.

© 2026 Abdurashidova Gulnura Abdurashidovna, Gulamov Ilhomjon Komiljonovich, & Haydarova Dilrabo Rahimjonovna. This work is licensed under a Creative Commons Attribution 4.0 International License (CC BY 4.0). The authors retain copyright and allow others to share, adapt, or redistribute the work with proper attribution.

**Cite This Article:** Abdurashidova Gulnura Abdurashidovna, Gulamov Ilhomjon Komiljonovich, & Haydarova Dilrabo Rahimjonovna. (2026). Results Of Evaluation Of Anti-Radical Properties Of *Inula Grandis Schrenk* (Asteraceae) Plant By DPPH Method. The American Journal of Agriculture and Biomedical Engineering, 8(01), 22–28. <https://doi.org/10.37547/tajabe/Volume08Issue01-04>

### 1. Introduction

Uzbekistan is a region naturally and geographically rich in medicinal plants, and of the approximately 4,500 species of higher plants naturally occurring in the republic, about 1,200 have medicinal properties [4,5]. The Resolution of the President of the Republic of Uzbekistan No. PQ-4670 dated April 10, 2020 “On measures for the protection, cultivation, processing and rational use of available resources of wild medicinal plants” created the legal basis for the radical development of the sector [17]

As a medicinal plant, *Inula grandis* Schrenk (Asteraceae) is a perennial plant. It grows on fine-grained soft slopes in the lower and middle mountain belts. The optimal resource potential is located at an altitude of 1300-1500 meters above sea level. According to modern indicators of resource potential in southern Uzbekistan, the permissible annual volume of raw material procurement is 5.16 tons [15]. Today, the phytochemistry of this species and its importance in folk medicine are being widely studied [16].


*Inula grandis* Schrenk is distributed in the South Chotkol region of the Fergana Valley (Namangan region, Yangikurgan district, Ungor-Tepa mountain) (Figure 1).

Human impact on the environment has increased significantly in the Fergana Valley over the past decade. The expansion of settlements, the growth of industrial zones, the reduction of pasture areas, and other factors have led to the weakening of the population of rare and endemic species of the Fergana Valley [1, 2, 3]. These indicators have increased the need to identify medicinal plant reserves in the regions and comprehensively study their phytochemistry.

In order to carry out research and analysis, the following tasks were set: 1) To comprehensively evaluate the antiradical activity of alcoholic and aqueous extracts

prepared from the *Inula grandis* plant using the DPPH method; 2) To determine the dependence of the antiradical activity of alcoholic and aqueous extracts on time and concentration, and to calculate their  $IC_{50}$  values based on linear regression models; 3) To experimentally substantiate the fact that the aqueous extract of *Inula grandis* has a higher antiradical potential than the alcoholic extract.

Funded by the Government of the Republic of Uzbekistan, and the project Development of the Digital Platform of the Flora of the Ferghana Valley as a 'Nature Imperilled Area' (Project No. AL-9224104319).



**Figure 1. A) Distribution map of *Inula grandis*; B) Ecology; C-D-E) Images of the plant in nature.**

## 2. Methods

Targeted and route-based field studies were conducted throughout the region during 2025 (May-August). The location coordinates of the species were determined using a GPS navigator (Garmin GPSmap 64s navigator). To obtain herbarium data, the TASH, MW (<https://www.plant.depo.msu.ru>) funds and the Global Biodiversity Information Facility (GBIF; <https://www.gbif.org>) were used. The nomenclature of taxa was given according to the international catalog

Plants of the World Online (POWO; <https://www.plantsoftheworldonline.org>) [6, 7, 8].

Laboratory analyses were carried out at Namangan State University. The discoloration of the purple 2,2-diphenyl-1-picrylhydrazyl (DPPH) solution allows us to determine the presence of some pure antioxidant compounds that have the properties of donating a hydrogen atom or an electron. Stable DPPH is a reagent used in spectrophotometric analysis. In this experiment, the method developed by Blois was used with minor

modifications to evaluate the inhibition of DPPH free radicals.

**Preparation of DPPH working solution.** A 7.92 mM DPPH solution in ethanol was prepared in a 100 ml volumetric flask, wrapped in aluminum foil, and stored in the dark at room temperature for 30 minutes [9, 10, 11].

**Preparation of sample extracts.** Alcoholic and aqueous extracts were prepared based on the selected sample. The preparation of sample extracts was carried out by extracting 1 g of plant sample in 25 ml of 96% ethanol and distilled water for 20 minutes in an ultrasonic bath. The obtained extract was passed through a 0.45 µm syringe filter and used for analysis. The filtered sample was diluted 10 times [12, 13, 14].

Determination of the antiradical properties of the samples. A 4 ml quartz cuvette was filled with 3 ml of DPPH solution and 100 µl of ethanol (blank sample) and placed in a spectrophotometer, and the absorbance (D1) at a wavelength of 517 nm was measured every 5 minutes for 30 minutes using a K7000 spectrophotometer manufactured by YOKE (China). To evaluate the antiradical properties of the sample, 25, 50, 75, 100 µl of the sample was mixed with 3 ml of DPPH solution and

the absorbance (D2) at 517 nm was measured in the above order. Ethanol was added to the remaining part to bring the total volume of the solution in the cuvette to 3.1 ml. The antiradical properties of the samples were calculated using the following formula:

$$\text{ARF\%} = \frac{(D_1 - D_2)}{D_1} \times 100\%$$


### 3. Results

100 µl of alcoholic extract inhibited DPPH free radicals by 31.47% in 30 minutes, while this figure was 50.73% for the aqueous extract. The IC<sub>50</sub> values calculated based on linear regression equations were 166.6 µl for the alcoholic extract and 100.1 µl for the aqueous extract, respectively. These results clearly showed that the aqueous extract had higher antiradical activity than the alcoholic extract

The results obtained suggest that the composition of *Inula grandis* is dominated by water-soluble phenolic and polyphenolic biologically active compounds. In general, the results of the study support the prospects for using *Inula grandis* as a source of natural antioxidants in the production of pharmaceutical, food and functional products (Figure 2).

**Table 1. Measured light absorption and antiradical activity of blank and test *Inula grandis* alcoholic extract added to DPPH solution**


Size, ml	Time minutes.	Sample					
		Abs, D <sub>2</sub>	ARF%		Time, minutes.	Abs, D <sub>2</sub>	ARF%
25	0	1,02	0,00	75	0	1,02	0,00
	5	0,952	6,67		5	0,885	13,24
	10	0,945	5,34		10	0,861	15,59
	15	0,938	5,92		15	0,848	16,86
	20	0,931	6,41		20	0,832	18,43
	25	0,926	6,80		25	0,821	19,51
	30	0,921	7,18		30	0,814	20,20
50	0	1,02	0,00	100	0	1,02	0,00
	5	0,936	8,24		5	0,798	21,76
	10	0,928	9,02		10	0,771	24,41



**Figure 2. Graphical representation of the measured light absorption of blank and tested alcohol-extracted *Inula grandis* solutions spiked with DPPH solution (Inula grandis alcoholic 10x).**

To calculate the *Inula grandis* IC50 – 50% inhibition concentration of DPPH solution, the following graph was constructed based on the 30-minute antiradical

scavenging activity (ARF%) values and the volume of alcohol samples added in each experiment and calculated based on the trend line function applied to it (Figure 3).




**Figure 3. Relationship graph between ARF% and volumes determined at 30 minutes of alcohol-extracted *Inula grandis* (Inula grandis alcoholic 10x)**

The trend line plotted on the graph was calculated from the functional formula  $y=mx+b$ , based on the formula  $x=(y-b)/m$ , which represents the volume that exhibits 50% ARF% (IC50):

$$IC_{50} = \frac{(50 - 0.410)}{0.298} = 166,6 \text{ mkl}$$

**Table 2. Measured light absorption and calculated antiradical activity values of blank and test aqueous extracts of *Inula grandis* added to DPPH solution**


size mkl	Time minutes.	Sample					
		Abs, D	ARF%		Time minutes.	Abs, D	ARF%
25	0	1,10	0,00	75	0	1,10	1,10
	5	0,996	9,45		5	0,932	0,996
	10	0,955	13,18		10	0,871	0,955
	15	0,924	16,00		15	0,829	0,924
	20	0,902	18,00		20	0,792	0,902
	25	0,885	19,55		25	0,764	0,885
	30	0,869	21,00		30	0,739	0,869
50	0	1,10	0,00	100	0	0	1,10
	5	0,961	12,64		5	5	0,711
	10	0,913	17,00		10	10	0,662
	15	0,878	20,18		15	15	0,621
	20	0,842	23,45		20	20	0,588
	25	0,819	25,55		25	25	0,563
	30	0,796	27,64		30	30	0,542



**Figure 4. Graphical representation of the measured light absorption of blank and tested aqueous extracted Inula grandis solutions spiked with DPPH solution (Inula grandis watery 10x).**

To calculate the IC<sub>50</sub> of Inula grandis - the 50% inhibitory concentration of DPPH solution, the following graph was constructed based on the 30-minute antiradical

scavenging activity (ARF%) values and the volume of alcohol samples added in each experiment and calculated based on the trend line function applied to it.



**Figure 5. Relationship graph between ARF% and volume determined at 30 minutes of aqueous extracted Inula grandis.**

The trend line plotted on the graph was calculated from the functional formula  $y=mx+b$ , based on the formula  $x=(y-b)/m$ , which represents the volume that exhibits 50% ARF% (IC<sub>50</sub>):

$$IC_{50} = \frac{(50 - 1.12)}{0.487} = 100,1 \text{ mkl}$$

**Table 3. Antiradical scavenging activity values (ARF%) of *Inula grandis* alcoholic extracts in 100 µl at 30 minutes**

	ARF%	
	Sample is alcohol	Sample is aqueous
30-minutes	31.47	50.73

**Table 4. IC50 of *Inula grandis* – 50% inhibition concentration of DPPH solution (µl)**

	ARF%	
	Sample is alcohol	Sample is aqueous
30-minutes	166.6	100.1

#### 4. Conclusion

As a result of the conducted studies, it was found that alcoholic and aqueous extracts prepared from the *Inula grandis* plant have significant antiradical activity against DPPH free radicals. Spectrophotometric analyses based on the DPPH method showed a consistent increase in antiradical activity with increasing extract volume and incubation time. Based on the results obtained, the scientific and practical significance of the *Inula grandis* plant as a natural antioxidant source is demonstrated.

#### Reference

1. Gulomov R.K. Distribution of the genus *Phlomoides* Moench in the Fergana Valley (taxonomy, geography, ecology and conservation): author's abstract of scientific paper, PhD – Tashkent: 2022– 43 p

2. Gulomov R.K., Batoshev A.R. Morphological Phylogeny of the Species *Phlomoides Moench* (Lamiaceae) Distributed in the Fergana Valley // International Journal of Virology and Molecular Biology. – USA, 2022. – № 11 (1). – P. 9-15. <https://doi.org/10.5923/j.ijvmb.20221101.03>

3. Gulomov, R., Rakhimova, K., Batoshev, A., Komilov, D. (2023). Bioclimatic modeling of the species *Phlomoides canescens* (Lamiaceae). The Scientific Temper, 14(4):1060-1065. <https://doi.10.58414/SCIENTIFICTEMPER.2023.14.4.01>.

4. Rahimova H.R., Gulomov R.K., Ibragimov A. DETERMINATION OF AMINO ACID AND VITAMIN CONTENT OF *Phlomoides nuda* AND *Phlomoides speciosa* (LAMIACEAE) SPECIES // Universum: химия и биология: электрон. научн. журн. 2023. 4(106). URL: <https://7universum.com/ru/nature/archive/item/15224>.

5. Khaydarova D.R. Lamiaceae obtaining medicinal food additives from certain types of herbs. author's abstract of scientific paper, PhD. - Andijan. 2022. – 46 p.

6. Moscow university virtual herbarium (<https://plant.depo.msu.ru>, accessed on 25 Aprel 2025)

7. Global Biodiversity Information Facility. 2025. Free and open acces to biodiversity data. Retrieved aprel 30, 2025, available from <https://www.gbif.org>.

8. Plants of the World Online. 2024. Facilitated by the Royal Botanic Gardens, Kew. Retrieved Aprel 30, 2025, available from <https://www.plantsoftheworldonline.org/>

9. Blois, M. S. (1958). Antioxidant determinations by the use of a stable free radical. Nature, 181, 1199–1200.

10. Gülcin, İ., Beydemir, Ş., Sat, İ. G., & Küfrevoiğlu, Ö. İ. (2005). Evaluation of antioxidant activity of cornelian cherry (*Cornus mas* L.). Acta Alimentaria Hungarica, 34, 193–202.

11. Askarov, I. R., Muminov, M. M., & Yusupov, M. A. (2024). Study of antiradical properties of artichoke (*Cynara scolymus* L.) and milk thistle (*Silybum marianum* L.) vegetable oils. NamDU Ilmiy Axborotnomasi, 11, 173–177.

12. Askarov, I. R., Abdullaev, S. S., Mamatkulova, S. A., & Abdulloev, O. S. (2024). Antioxidant activity

and elemental composition of mixtures of fig and common unabi fruits. *Journal of Chemistry of Goods and Traditional Medicine*, 3(3), 179–205.  
<https://doi.org/10.55475/jcgtn/vol3/iss3/2024/320>

13. Prior, R. L., Wu, X., & Schaich, K. (2005). Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *Journal of Agricultural and Food Chemistry*, 53(10), 4290–4302.
14. Brand-Williams, W., Cuvelier, M. E., & Berset, C. (1995). Use of a free radical method to evaluate antioxidant activity. *LWT - Food Science and Technology*, 28(1), 25–30.
15. Mahmudov A.V., Abduraimov O.S., Allamurotov A.L., Maxmudov V., Mavlanov B.J. & Mamatkasimov O.T., "Resources of Inula grandis Schrenk ex Fisch. & C.A. Mey. in South Uzbekistan," *Plant Science Today*, vol. 11, no. 3, pp. 719–727, 2024.  
<https://doi.org/10.14719/pst.3392>.
16. Azizbek Maximudov, Ozodbek Abduraimov, Akmal Allamurotov, Valijon Maximudov, Shoxina Halimova, Komiljon Tojibayev, "Scale for Assessing the Impact of Operational Stress on the Resource Potential of Medicinal Plants: A Case Study of Southern Uzbekistan," *Environment and Ecology Research*, Vol. 13, No. 6, pp. 850 - 867, 2025. DOI: 10.13189/eer.2025.130608.
17. Resolution of the President of the Republic of Uzbekistan No. PQ-4670 dated April 10, 2020 "On measures for the protection, cultivation, processing and rational use of available resources of wild medicinal plants"