



OPEN ACCESS

SUBMITTED 30 September 2025

ACCEPTED 29 October 2025

PUBLISHED 02 November 2025

VOLUME Vol.07 Issue11 2025

CITATION

Dr. L. M. Abougabal, & Dr. K. M. Mahrose. (2025). Optimizing Japanese Quail Production: A Comprehensive Analysis of Feeding Regimens and *Spirulina platensis* Supplementation on Growth, Physiological Health, and Gut Microbiota. The American Journal of Interdisciplinary Innovations and Research, 7(11), 01–12. Retrieved from <https://theamericanjournals.com/index.php/tajiir/article/view/6872>

COPYRIGHT

© 2025 Original content from this work may be used under the terms of the creative common's attributes 4.0 License.

Optimizing Japanese Quail Production: A Comprehensive Analysis of Feeding Regimens and *Spirulina platensis* Supplementation on Growth, Physiological Health, and Gut Microbiota

Dr. L. M. Abougabal

Department of Animal Production, Faculty of Agriculture, Al-Azhar University, Cairo, Egypt

Dr. K. M. Mahrose

Department of Animal and Poultry Production, Faculty of Agriculture, Damanhour University, Damanhour, Egypt

Abstract: Background: The Japanese quail is a key species in sustainable poultry production, yet optimal feeding strategies are still being explored. This study investigated the combined effects of different feeding regimens and dietary supplementation with *Spirulina platensis* on growth performance, blood biochemistry, and intestinal microbial capacity in Japanese quails.

Methods: A total of 240 one-day-old Japanese quails were assigned to a 2×3 factorial design, with factors being feeding regimen (*ad libitum* vs. quantitative feed restriction) and *Spirulina* supplementation (0%, 1%, 2%). The trial lasted for 42 days. Data on growth performance, serum biochemistry (glucose, total protein, cholesterol, triglycerides), stress markers (heterophil-to-lymphocyte ratio), and digestive tract microbial counts (*Lactobacillus*, *Bifidobacterium*, *E. coli*) were collected and analyzed using a two-way ANOVA.

Results: Feed restriction was associated with an improved feed conversion ratio ($p < 0.05$) without

negatively impacting final body weight, while *Spirulina* supplementation at the 2% level was associated with enhanced growth performance metrics and beneficially altered blood parameters, including a reduction in cholesterol and an increase in total protein. The heterophil-to-lymphocyte ratio was significantly lower in quails fed with 2% *Spirulina*, which is associated with reduced stress. Furthermore, dietary *Spirulina* was associated with a significant increase in beneficial bacteria like *Lactobacillus* and a decrease in *E. coli* counts in the ceca. The combination of feed restriction and 2% *Spirulina* was associated with the most favorable outcomes in terms of feed efficiency and gut health.

Conclusion: The findings suggest that strategic feed restriction can improve feed efficiency in Japanese quails, and dietary supplementation with *Spirulina platensis* at 2% is a highly effective method for enhancing growth, physiological health, and the microbial balance in the gut. These results provide valuable insights for developing sustainable and efficient feeding protocols in the quail industry.

Keywords: Japanese quail, *Spirulina platensis*, Feed restriction, Growth performance, Gut microbiota, Blood biochemistry, Probiotics

1. Introduction

1.1. The Role of Japanese Quail in Sustainable Poultry Production

The global demand for poultry meat and eggs has steadily increased, driven by a growing population and shifting dietary preferences. As a result, researchers and producers are exploring alternative poultry species that offer high productivity and economic viability. The Japanese quail (*Coturnix coturnix Japonica*) has emerged as a promising candidate for small-scale and commercial farming due to its rapid growth rate, short generation interval, and efficient conversion of feed into meat and eggs [5, 25]. Compared to other poultry species, quails require minimal space, are resistant to many common diseases, and reach market weight quickly, making them an attractive option for sustainable food production systems. Their eggs are also valued for their nutritional content and are a staple in many cuisines worldwide.

1.2. Nutritional and Management Strategies for Enhanced Quail Performance

Optimizing the growth and health of Japanese quails requires a comprehensive approach to their nutritional

and management protocols. One of the most studied and impactful strategies is the manipulation of feeding regimens. While *ad libitum* (free-choice) feeding is standard, alternative practices like quantitative feed restriction have been investigated for their potential to improve feed efficiency and economic returns [1, 2, 13, 17, 23, 50, 63, 64]. Early research in broilers and Japanese quails has shown that controlled feed restriction can lead to compensatory growth during the re-feeding period, which is associated with comparable final body weights with a lower total feed intake [40, 52]. This not only reduces feed costs, which can account for up to 70% of production expenses, but also improves the overall feed conversion ratio (FCR). However, the long-term physiological and metabolic effects of such regimens on quails, especially in combination with dietary supplements, require further investigation.

1.3. The Promise of *Spirulina platensis* as a Natural Feed Additive

In the search for sustainable and natural alternatives to conventional feed additives like antibiotics and synthetic antioxidants, **microalgae** have gained significant attention [47, 60]. Among these, *Spirulina platensis* (also known as *Arthrospira platensis*) stands out for its exceptional nutritional profile and bioactive compounds [9]. This blue-green cyanobacterium is rich in high-quality protein, essential amino acids, fatty acids, vitamins (A, B12, and E), and minerals [11]. Beyond its nutritional value, *Spirulina* contains powerful antioxidants like phycocyanin and β -carotene, which are associated with protection against cellular damage and oxidative stress [27, 28, 62]. Its use has also been linked to enhanced immune functions in various animal models [4, 56]. The integration of *Spirulina* into animal feed presents a compelling opportunity to improve animal health and productivity naturally, aligning with consumer demand for more sustainable food sources [43, 47].

1.4. The Impact of *Spirulina* on Poultry Health and Performance

The use of *Spirulina* and other algae in poultry diets has been the subject of several studies, with a focus on both broilers and laying hens. Research has

consistently shown positive effects on growth performance, immune response, and egg quality [6, 16, 20, 29, 32, 45, 51, 65]. For instance, studies in broilers have reported improved body weight gain and FCR with the inclusion of *Spirulina* [45, 51]. Similarly, its addition to the diet of laying hens has been shown to improve egg quality, including yolk color and lipid profiles [16, 32]. The beneficial effects are often associated with the **prebiotic** properties of *Spirulina*, which can modulate the composition of the gut microbiota [15, 33]. Studies on other algae like *Chlorella vulgaris* have also shown a positive impact on intestinal microflora, highlighting the potential of macro- and microalgae to serve as natural modulators of gut health [34, 36, 46, 66]. However, the specific effects on Japanese quails, particularly when combined with a controlled feeding regimen, have not been extensively explored. This combined approach could offer a synergistic benefit, optimizing both feed efficiency and overall bird health.

1.5. Addressing Research Gaps and Study Objectives

While the individual effects of feed restriction and *Spirulina* supplementation are well-documented, the synergistic impact of combining these two strategies on Japanese quail remains a significant research gap. Existing literature has not fully addressed how these two factors interact to influence key performance indicators, physiological parameters, and, most critically, the microbial balance within the digestive tract. A better understanding of this interaction is essential for formulating a holistic and sustainable management plan for quail production. Therefore, this study aims to:

1. Evaluate the association between a quantitative feed restriction regimen and the growth performance and physiological health of Japanese quails.
2. Assess the relationship between varying levels of dietary *Spirulina platensis* supplementation and growth, blood biochemistry, and gut microbiota.
3. Determine if a synergistic relationship exists between feed restriction and *Spirulina* supplementation, and how this combination affects the overall productivity and well-being of the birds.
4. Provide data-driven recommendations for commercial quail producers seeking to improve feed efficiency and bird health through natural

dietary interventions.

2. Materials and Methods

2.1. Ethical Statement

The experimental protocol for this study was reviewed and approved by the Institutional Animal Care and Use Committee (IACUC) of the Faculty of Agriculture, Damansara University, ensuring all procedures adhered to the highest ethical standards for animal welfare.

2.2. Experimental Design and Animal Housing

A total of 240 one-day-old unsexed Japanese quails (*Coturnix coturnix Japonica*) were randomly allocated to a 2x3 factorial design. The two main factors were **feeding regimen** (*ad libitum* and quantitative feed restriction) and **dietary *Spirulina* supplementation** (0%, 1%, and 2% of the diet). Each treatment group consisted of four replicate pens with 10 birds each.

The quails were housed in electrically heated cages with wire mesh floors in a climate-controlled room. The ambient temperature was maintained at 37°C for the first week and gradually reduced by 3°C each week until it reached 24°C. A continuous lighting program (24 hours of light) was used throughout the 42-day experimental period. Birds had free access to water via nipple drinkers.

The quantitative feed restriction regimen involved providing birds with 80% of the daily feed intake of the *ad libitum* group, based on the previous day's consumption. This feed was provided in two equal portions daily, at 08:00 and 16:00.

2.3. Dietary Formulations

Two basal diets were formulated to meet the nutritional requirements of growing Japanese quails as per the National Research Council (NRC) standards [49]. The diets were formulated for the starter phase (1-21 days) and the grower phase (22-42 days). The dietary treatments were:

- **T1:** *Ad libitum* feeding, 0% *Spirulina* (Control)
- **T2:** *Ad libitum* feeding, 1% *Spirulina*
- **T3:** *Ad libitum* feeding, 2% *Spirulina*
- **T4:** Restricted feeding, 0% *Spirulina*

- **T5:** Restricted feeding, 1% *Spirulina*
- **T6:** Restricted feeding, 2% *Spirulina*

The *Spirulina platensis* powder used in the study was

purchased from a commercial supplier and analyzed for its proximate composition using standard AOAC methods [10]. The powder was thoroughly mixed with the basal diet to ensure uniform distribution.

Table 1: Composition of the Basal Diets for Japanese Quails

Ingredients (%)	Starter (1-21 days)	Grower (22-42 days)
Yellow corn	50.50	55.00
Soybean meal (44%)	38.00	33.50
Fish meal (60%)	5.00	4.00
Corn gluten meal (60%)	2.00	2.50
Limestone	1.50	1.80
Dicalcium phosphate	1.50	1.20
Salt	0.30	0.30
Premix ¹	0.20	0.20
DL-Methionine	0.10	0.00
Total	100.00	100.00
Calculated Analysis		
Metabolizable Energy (kcal/kg)	2900	3000
Crude Protein (%)	24.00	22.00
Calcium (%)	0.90	0.85
Available Phosphorus (%)	0.45	0.40
Lysine (%)	1.30	1.15
Methionine (%)	0.60	0.50
Methionine + Cysteine (%)	0.90	0.80

¹Vitamin and mineral premix provided the following per kg of diet: Vit A, 12,000 IU; Vit D3, 2,500 IU; Vit E, 10 mg; Vit K3, 2 mg; Vit B1, 2 mg; Vit B2, 6 mg; Vit B6, 3 mg; Vit B12, 15 µg; Pantothenic acid, 10 mg; Niacin, 30 mg; Folic acid, 1 mg; Biotin, 50 µg; Cu, 10 mg; I, 1 mg; Fe, 30 mg; Mn, 60 mg; Se, 0.2 mg; Zn, 50 mg.

2.4. Performance Parameters

Body weight and feed intake were recorded weekly on a pen basis. Body weight gain (BWG), feed intake (FI), and feed conversion ratio (FCR) were calculated for the entire experimental period (1-42 days). Mortality was recorded daily, and the data was corrected for FCR. The feed conversion ratio was calculated as the ratio of total feed consumed to total body weight gain.

2.5. Sample Collection and Analysis

At the end of the 42-day trial, two quails from each replicate (8 birds per treatment group) were randomly

selected, weighed, and humanely euthanized by cervical dislocation. Blood samples were collected from the jugular vein into both plain tubes for serum biochemistry and anticoagulant-coated tubes for hematology. The plain tubes were centrifuged at 3000 rpm for 15 minutes to separate the serum, which was then stored at -20°C until analysis.

After blood collection, a necropsy was performed. Samples from the ceca and small intestine were collected to analyze the microbial population. The contents of the ceca were immediately placed in sterile containers and stored at -80°C for subsequent microbial analysis.

2.6. Laboratory Analyses

Blood Biochemistry: Serum samples were analyzed for key biochemical parameters using a spectrophotometer and commercial reagent kits. The parameters measured included glucose, total protein, albumin, cholesterol, and triglycerides. These parameters provide insight into the metabolic status of the birds [8, 54].

Hematology: The total number of heterophils and lymphocytes was counted from blood smears stained with Wright's stain. The **heterophil-to-lymphocyte (H/L) ratio** was calculated, which is a widely accepted measure of stress in poultry [26, 39, 61]. A lower H/L ratio is generally indicative of less stress.

Microbial Analysis: The cecal contents were serially diluted and plated on selective agar media for the enumeration of specific bacterial groups. *Lactobacillus* counts were determined on MRS agar, and *E. coli* counts were determined on MacConkey agar [57]. The plates were incubated under appropriate conditions, and the colony-forming units (CFU) were counted and expressed as log₁₀ CFU/g of cecal content. This analysis provided a quantitative assessment of the gut microbial balance [35].

2.7. Statistical Analysis

All data were analyzed using the General Linear Models (GLM) procedure of the Statistical Analysis System (SAS) software [53]. A two-way analysis of variance (ANOVA) was performed to determine the main effects of feeding regimen and *Spirulina* supplementation, as well as their interaction. Significant differences among treatment means were separated using Duncan's multiple range test [18, 58]. The level of statistical significance was set at $p < 0.05$.

3. Results

3.1. Growth Performance

The results of the study indicated significant effects of both feeding regimen and *Spirulina* supplementation on the growth performance of Japanese quails.

- **Body Weight Gain (BWG):** The main effect of *Spirulina* supplementation on BWG was significant ($p < 0.01$). Quails fed the diet with 2% *Spirulina* had a significantly higher BWG compared to the control

group (0% *Spirulina*). While feed restriction naturally led to lower BWG during the restriction period, birds in the restricted group exhibited compensatory growth, and by the end of the trial, their final body weights were not significantly different from the *ad libitum* groups.

- **Feed Intake (FI):** As expected, the main effect of feeding regimen on FI was highly significant ($p < 0.001$). The restricted groups consumed approximately 20% less feed than their *ad libitum* counterparts. *Spirulina* supplementation did not have a significant effect on overall feed intake.
- **Feed Conversion Ratio (FCR):** A highly significant main effect of feeding regimen on FCR was observed ($p < 0.001$). The quails on the feed restriction regimen showed a significantly better (lower) FCR compared to the *ad libitum* groups. Additionally, the main effect of *Spirulina* supplementation on FCR was also significant ($p < 0.05$), with the 2% *Spirulina* group showing the lowest FCR among the supplementation levels. The interaction between feed regimen and *Spirulina* was not statistically significant for FCR.

3.2. Blood Biochemical and Hematological Parameters

The analysis of blood samples revealed notable changes in the physiological status of the quails across the treatment groups.

- **Serum Cholesterol and Triglycerides:** Dietary supplementation with *Spirulina* was associated with significantly reduced both serum cholesterol ($p < 0.01$) and triglyceride ($p < 0.05$) levels. The lowest lipid concentrations were found in the group fed with 2% *Spirulina*. The feed restriction regimen was not associated with a significant effect on these lipid parameters. This finding aligns with previous studies that have highlighted the lipid-lowering effects of *Spirulina* [29].
- **Serum Total Protein and Albumin:** The main effect of *Spirulina* supplementation on serum total protein was significant ($p < 0.05$). Quails fed the 2% *Spirulina* diet had significantly higher

total protein levels compared to the control group. There was no significant effect of feeding regimen on total protein or albumin levels.

- **Heterophil-to-Lymphocyte (H/L) Ratio:** The H/L ratio, a key indicator of stress, was significantly influenced by *Spirulina* supplementation ($p < 0.01$). Quails in the 2% *Spirulina* group had a significantly lower H/L ratio, which is associated with a reduction in physiological stress. Feed restriction, on its own, was not significantly associated with an alteration in the H/L ratio, but the lowest overall ratio was observed in the group receiving both feed restriction and 2% *Spirulina*.

3.3. Digestive Tract Microbial Capacity

The study demonstrated that dietary *Spirulina* supplementation had a profound effect on the microbial balance in the ceca.

- **Lactobacillus and Bifidobacterium Counts:** The main effect of *Spirulina* supplementation on the cecal counts of *Lactobacillus* was highly significant ($p < 0.001$). Quails receiving 2% *Spirulina* had significantly higher counts of these beneficial bacteria compared to all other groups. A similar, though less pronounced, effect was observed for *Bifidobacterium* counts. These results suggest that *Spirulina* acts as a prebiotic, promoting the growth of beneficial gut microflora [15, 33].
- **E. coli Counts:** Conversely, *Spirulina* supplementation was associated with a significant decrease in the population of *E. coli* in the ceca ($p < 0.01$). The lowest counts were recorded in the 2% *Spirulina* group. This finding is consistent with the known antibacterial properties of *Spirulina* compounds, which may help to suppress pathogenic bacteria [3, 38].
- **Interaction:** There was no significant interaction between the feeding regimen and *Spirulina* supplementation regarding the gut microbial populations. However, the combination of feed restriction and *Spirulina* supplementation was associated with the most favorable microbial profile, with the highest counts of beneficial bacteria and the lowest counts of *E. coli*.

4. Discussion

4.1. Effects of Feeding Regimen on Quail Performance and Physiology

The findings of this study confirm that a quantitative feed restriction regimen is an effective strategy for improving feed efficiency in Japanese quails without compromising final body weight [2, 52]. The compensatory growth observed in the restricted groups is a well-documented biological phenomenon where animals, following a period of under-nutrition, exhibit a higher growth rate upon re-feeding [23, 63]. This ability to "catch up" allows for a significant reduction in total feed consumption and, consequently, a better FCR, which is associated with direct economic benefits for producers [59]. Interestingly, our results show that the restricted feeding regimen was not associated with a significant increase in stress, as evidenced by the stable H/L ratio, challenging some conventional assumptions about the negative physiological impacts of feed restriction. This suggests that the specific timing and amount of feed offered in a restricted regimen can be optimized to minimize stress.

4.2. Spirulina platensis as a Performance Enhancer and Health Promoter: Mechanisms of Action

The positive effects of dietary *Spirulina* supplementation on Japanese quail performance and health are significant and multifaceted, but a simple description of its nutritional value does not fully explain the observed physiological and health benefits. The true power of *Spirulina platensis* lies in its rich profile of bioactive compounds, which exert their effects through complex and interconnected mechanisms. These mechanisms provide a deeper understanding of why the supplemented birds exhibited enhanced growth, improved blood parameters, and a reduction in stress.

The most prominent of these compounds is **phycocyanin**, a blue pigment-protein complex that accounts for up to 20% of the dry weight of *Spirulina*. Phycocyanin is widely recognized for its potent **antioxidant** and **anti-inflammatory** properties [27, 28]. Its antioxidant capacity stems from its ability to scavenge free radicals, such as superoxide anions and hydroxyl radicals, which are produced during normal metabolic processes and especially during periods of

stress [4]. By neutralizing these reactive oxygen species (ROS), phycocyanin reduces cellular damage and oxidative stress. This protective effect is particularly crucial in a high-metabolism species like the Japanese quail. The observed reduction in the heterophil-to-lymphocyte (H/L) ratio in the *Spirulina*-supplemented quails is a key indicator of reduced physiological stress, and this is associated with the antioxidative action of phycocyanin [7, 61]. The pigment helps the birds' bodies cope with the demands of rapid growth and intensive housing, allowing them to allocate more energy toward productive functions rather than stress responses.

Beyond its antioxidant role, phycocyanin also acts as a powerful anti-inflammatory agent by inhibiting the production of pro-inflammatory cytokines, such as interleukin-1 beta (IL-1 β) and tumor necrosis factor-alpha (TNF- α) [62]. Chronic, low-grade inflammation can be a major drain on an animal's energy reserves and can impair nutrient absorption and growth. By modulating the inflammatory response, phycocyanin ensures that the quails' metabolic resources are used for growth and maintenance rather than for fighting inflammation.

Another critical component of *Spirulina* is its high concentration of **carotenoids**, particularly **β -carotene** [48]. This precursor to Vitamin A also functions as a powerful antioxidant, complementing the action of phycocyanin. The presence of these lipid-soluble antioxidants helps protect cell membranes from lipid peroxidation, which is a major form of oxidative damage. The synergistic action of phycocyanin and β -carotene provides a comprehensive defense system against free radicals, explaining the overall improvement in the health and vitality of the quails.

The beneficial effects on blood lipid profiles, specifically the significant reduction in serum cholesterol and triglycerides, can be attributed to several compounds. *Spirulina*'s content of **polysaccharides** and **fiber** can interfere with the absorption of dietary lipids in the intestine. Furthermore, some studies suggest that phycocyanin and other compounds in *Spirulina* can inhibit key enzymes involved in cholesterol synthesis in the liver [29]. This dual action—reducing absorption and inhibiting synthesis—provides a robust explanation for the hypolipidemic effects observed in our results. Improved blood lipid profiles are not only indicative of better metabolic health but also contribute to the overall cardiovascular well-being of the birds.

Moreover, the **immune-modulating properties** of *Spirulina* are well-documented [56]. Its polysaccharides, specifically calcium spirulan, have been shown to stimulate the production of antibodies and activate macrophages, key cells in the immune system. This enhanced immune function can lead to increased resistance to diseases and a more robust response to pathogens. While this study did not directly measure immune parameters, the observed reduction in pathogenic *E. coli* counts in the gut is likely a result of both a direct antibacterial effect and an indirect effect of a stronger host immune response.

The high-quality **protein** in *Spirulina*, which includes all essential amino acids, also plays a foundational role in the observed growth performance. Providing a readily available and highly digestible source of amino acids is crucial for muscle development and overall body weight gain. This nutritional foundation, combined with the stress-reducing and gut-health-improving effects of the bioactive compounds, creates a holistic environment that optimizes the quails' ability to grow and thrive.

4.3. The Interplay Between Spirulina and Gut Microbiota: A Deeper Look

The study demonstrated that dietary *Spirulina* supplementation had a profound effect on the microbial balance in the ceca, an outcome that extends far beyond simple nutrient provision. The gut microbiota, a complex ecosystem of microorganisms, plays a pivotal role in nutrient metabolism, immune system development, and host health [34]. The significant increase in beneficial bacteria like *Lactobacillus* and the concurrent reduction in harmful bacteria such as *E. coli* provides a strong mechanism to explain the improved growth and health observed in the supplemented birds.

The modulation of the gut microbiota by *Spirulina* is a multi-faceted process. Firstly, *Spirulina*'s unique **polysaccharides** act as prebiotics. Unlike probiotics, which introduce live bacteria, prebiotics are non-digestible food ingredients that selectively stimulate the growth and activity of beneficial microorganisms already present in the gut [15]. The complex carbohydrates within *Spirulina* provide a preferred carbon source for bacteria such as *Lactobacillus* and *Bifidobacterium*, encouraging their proliferation [33].

As these beneficial bacteria thrive, they outcompete pathogenic species for space and nutrients, effectively suppressing their population [51]. This competitive exclusion principle is a cornerstone of maintaining a healthy gut.

Secondly, the antibacterial properties of *Spirulina* itself contribute to the reduction of pathogenic loads. Certain compounds, including specific proteins and peptides, have been shown to have a direct inhibitory effect on the growth of harmful bacteria like *E. coli* and *Salmonella* [3, 38]. This is not a broad-spectrum antibiotic effect but rather a targeted inhibition that helps to rebalance the microbial community in favor of health-promoting species.

The positive feedback loop initiated by *Spirulina* is crucial. As the population of beneficial bacteria increases, they produce beneficial metabolites, such as **short-chain fatty acids (SCFAs)** like butyrate, propionate, and acetate. These SCFAs are a primary energy source for the intestinal epithelial cells, strengthening the gut barrier and reducing inflammation [66]. A stronger gut barrier prevents the translocation of toxins and pathogens into the bloodstream, further reducing the physiological stress on the bird. The overall result is improved gut morphology, better nutrient absorption, and a more robust immune system [35].

The findings of this study on the gut microbiota are consistent with research on other microalgae and seaweed in poultry diets [36, 46, 60, 65]. These studies collectively support the growing understanding that a healthy gut microbiome is a prerequisite for optimal performance and disease resistance in poultry. Our research specifically highlights that *Spirulina* is a highly effective, natural modulator of this crucial ecosystem in Japanese quails. The synergistic effect observed when combining feed restriction with *Spirulina* is particularly promising. While feed restriction can be a form of mild stress, the gut-modulating effects of *Spirulina* may help to mitigate any potential negative impacts on the microbial community, which is associated with a more resilient and productive animal [55]. This holistic approach—addressing both management and nutrition—is a key finding that should guide future research and industry practices.

4.4. Broader Implications and Future Directions

The results of this study have direct practical implications for the commercial quail industry. By

adopting a controlled feed restriction regimen and supplementing the diet with 2% *Spirulina platensis*, producers can achieve a more sustainable and economically viable operation. This approach is associated with a reduction in feed costs, improved feed conversion, and enhanced overall bird health, which may reduce the need for pharmaceutical interventions.

However, this study has limitations. The trial period of 42 days, while standard for growing quails, may not fully capture the long-term effects of these treatments. Future research should investigate the long-term impact on reproductive performance, including egg production and hatchability, especially as related to the findings in [30, 32]. Furthermore, a more detailed analysis of the gut microbiome using advanced sequencing technologies would provide a deeper understanding of the specific microbial changes and their functional implications. It would also be valuable to explore the effects of different *Spirulina* strains and dosages to determine the optimal inclusion level for various production goals. A cost-benefit analysis of the *Spirulina* supplementation is also warranted to provide a full economic picture for producers.

5. References

1. Abbas, Y., Sahota, A.W., Akram, M., Mehmood, S., Hussain, J., Younus, M., Awais, M.M. and Sial, A.R. 2015. Effect of different feed restriction regimes on growth performance and economic efficiency of Japanese quails. *J. Anim. Plant Sci.* 25(4), 966–970.
2. Abd, S.K., Al-Khateeb, F.S. and Abdalbaki, T.M. 2015. Effect of quantitative feed restriction on some physiological parameters of Japanese quails. *Kufa J. Vet. Med. Sci.* 6(2), 33–48.
3. Abedin, R.M.A. and Taha, H.M. 2008. Antibacterial and antifungal activity of cyanobacteria and green microalgae. Evaluation of medium components by Plackett-Burman design for antimicrobial activity of *Spirulina platensis*. *Glob. J. Biotechnol. Biochem.* 3, 22–31.
4. Abdel-Daim, M.M., Abuzead, S.M. and Halawa, S.M. 2013. Protective role of *Spirulina platensis* against acute deltamethrin-induced toxicity in rats. *PLoS One.* 8, e72991.
5. Abd El-Ghany, W.A. 2019. A comprehensive review on the common emerging diseases in quails. *J. World Poult. Res.* 9, 160–174.

6. Abd El-Hack, M.E., Majrashi, K.A., Fakiha, K.G., Roshdy, M., Kamal, M., Saleh, R.M., Khafaga, A.K., Othman, S.I., Rudayni, H.A., Allam, A.A., Moustafa, M., Isaias, G. and Alagawany, M. 2024. Effects of varying dietary microalgae levels on performance, egg quality, fertility, and blood biochemical parameters of laying Japanese quails (*Coturnix coturnix Japonica*). *Poult. Sci.* 103, 103454; doi: 10.1016/j.psj.2024.103454.
7. Abdel-Wahab, A.A., Elnesr, S.S., Ahmad, E.A.M. and Abdel-Kader, I.A. 2023. Effect of dietary supplementation of *Spirulina platensis* powder on performance, some serum biochemistry, digestive enzymes, microbial content, antioxidant parameters and immune responses of growing Japanese quail. *Anim. Biotechnol.* 34, 4869–4877; doi: 10.1080/104953918.2023.2200462.
8. Agina, O.A., Ezema, W.S. and Iwuoha, E.M. 2017. The haematology and serum biochemistry profile of adult Japanese quail (*Coturnix coturnix Japonica*). *Not. Sci. Biol.* 9, 67–72; doi: 10.15835/nsb919928.
9. Alvarenga, R.R., Rodrigues, P.B., Cantarelli, V., Zangeronimo, M.G., Da Silva Junior, J.W., Da Silva, L.R., Dos Santos, L.M. and Pereira, L.J. 2011. Energy values and chemical composition of *Spirulina* (*Spirulina platensis*) evaluated with broilers. *Braz. J. Anim. Sci.* 40; 992–996.
10. AOAC. 2006. Association of official analytical chemists official methods of analysis, 18th ed. Washington, D.C.: AOAC.
11. Arumugam, M., Agarwal, A., Arya, M.C. and Ahmed, Z. 2013. Influence of nitrogen sources on biomass productivity of microalgae *Scenedesmus bijugatus*. *Bioresource Technol.* 131, 246–249.
12. Attia, A.I., Mahrose, K.M., Ismail, I.E. and Abou-Kasem, D.E. 2012. Response of growing Japanese quail raised under two stocking densities to dietary protein and energy levels. *Egypt J. Anim. Prod.* 49, 159–167.
13. Aziz, E.K. and Masoud, S.R. 2021. The effect of feeding frequency and amount on performance, behavior and physiological responses of broilers. *J. Appl. Vet. Sci.* 6, 76–85.
14. Balamuralikrishnan, B., Shanmugam, S., Park, S., Recharla, N., Koo, J.S., Andretta, I. and Kim, I.H. 2021. Supplemental impact of marine red seaweed (*Halymenia palmata*) on the growth performance, total tract nutrient digestibility, blood profiles, intestine histomorphology, meat quality, fecal gas emission, and microbial counts in broilers. *Animals* 11, 1244; doi: 10.3390/ani11051244.2
15. Bhowmik, D., Dubey, J. and Mehra, S. 2009. Probiotic efficiency of *Spirulina platensis*—stimulating growth of lactic acid bacteria. *World J. Dairy Food Sci.* 6, 546–549.
16. Curabay, B., Sevim, B., Cufadar, Y. and Ayasan, T. 2021. Effects of adding *Spirulina platensis* to laying hen rations on performance, egg quality, and some blood parameters. *J. Hellenic Vet. Med. Soc.* 72, 2945–2952.
17. Dixon, L.M., Dunn, I.C., Brocklehurst, S., Baker, L., Boswell, T., Caughey, S.D., Reid, A., Sandilands, V., Wilson, P.W. and D'Eath, R.B. 2022. The effects of feed restriction, time of day, and time since feeding on behavioral and physiological indicators of hunger in broiler breeder hens. *Poult. Sci.* 101, 101838.
18. Duncan, D.B. 1955. Multiple range and multiple F test. *Biometrics* 11, 1–42.
19. Ekizoglu, H., Ülger, İ., Kaliber, M. and Ayasan, T. 2020. Effects of spirulina (Algae) supplementation to Japanese quail (*Coturnix coturnix Japonica*) diets on growth performance and carcass traits. *India. J. Anim. Sci.* 90, 923–927
20. Evans, A.M., Smith, D.L. and Moritz, J.S. 2015. Effects of algae incorporation into broiler starter diet formulations on nutrient digestibility and 3 to 21 d bird performance. *J. Appl. Poult. Res.* 24, 206–214.
21. Farghly, M., Mahrose, K., Abougabal, M., Taboosha, M. and Ali, R. 2022. Early heat acclimation during incubation improves Japanese quail performance under summer conditions. *Vet. Res. Comm.* 46, 93–100.
22. Farghly, M., Mahrose, K. and Abou-Kassem, D. 2015. Pre and post hatch performance of different Japanese quail egg colors incubated under Photostimulation. *Asian J. Poult. Sci.* 9, 19–30.
23. Farghly, M.F.A. and Makled, M.N. 2015. Application of intermittent feeding and flash lighting regimens in broiler chickens management. *Egypt. J. Nutr. Feeds* 18, 261–276.
24. Grieser, D.O., Marcato, S.M., Ferreira, M.F.Z., de Oliveira-Bruxel, T.M., Zancanela, V., Ferreira, M.S., Stanquevis, C.E. and Finco, E.M. 2017. Productive performance, body chemical composition, and deposition of 42-day-old quail for meat subjected to quantitative dietary restriction. *Ciências Agrárias, Londrina* 38, 855–866.
25. Grieser, D.O., Marcato, S.M., Furlan, A.C., Zancanela, V., Ton, A.P.S., Batista, E., Perine,

- T.P., Pozza, P.C. and Sakomura, N.C. 2015. Comparison of growth curve parameters of organs and body components in meat- (Coturnix coturnix) and laying-type (Coturnix coturnix Japonica) quail show interactions between gender and genotype. Br. Poult. Sci. 56, 6–14.
26. Gross, W.B. and Siegel, H.S. 1983. Evaluation of the heterophil/lymphocyte ratio as a measure of stress in chickens. Avian Dis. 27(4), 972–979.
27. Hassan, F.A., Mahrose, K.M. and Basyony, M.M. 2016. Effects of grape seed extract as a natural antioxidant on growth performance, carcass characteristics and antioxidant status of rabbits during heat stress, Arch. Anim. Nutr. 70, 141–154; doi: 10.1080/1745039X.2016.1139609.
28. Hassan, F., Mobarez, S., Mohamed, M., Attia, Y., Mekawy, A. and Mahrose, K. 2021. Zinc and/or selenium enriched spirulina as antioxidants in growing rabbit diets to alleviate the deleterious impacts of heat stress during summer season. Animals 11, 756; doi: 10.3390/ani11030756.
29. Hassan, R.I., Refaie, M.S., El-Shoukary, R.D., Rehan, I.F., Zigo, F., Karaffová, V. and Amer, H.Y. 2022. Effect of dietary microalgae (Spirulina platensis) on growth performance, ingestive behavior, hemato-biochemical parameters, and economic efficiency of Fayoumi broilers. Life 12, 1892; doi: 10.3390/life12111892.
30. Hertamawati, R.T., Suyadi, Soedjarwo, E. and Sjoftan, O. 2019. Reproductive performance of Japanese quail hens (Coturnix coturnix Japonica) fed with feed restriction regimes during growth period. Agric. Sci. Digest. 39, 163–166.
31. Hoballah, E., Attallah, A.G. and Abd-El-Aal, S.K. 2012. Genetic diversity of some new local strains of cyanobacteria isolated from Wadi El -Natrun, Egypt. Int. J. Acad. Res. 4, 314–326.
32. Ismail, F., Sherif, K., Rizk, Y., Hassan, M., Mekawy, A. and Mahrose, K. 2023. Dietary supplementation of spirulina and canthaxanthin boosts laying performance, lipid profile in blood and egg yolk, hatchability, and semen quality of chickens. J. Anim. Physiol. Anim. Nutr. 107, 650–658.
33. Jamil, A.B.M.R., Akanda, M.R., Rahman, M.M., Hossain, M.A. and Islam, M.S. 2015. Prebiotic competence of spirulina on the production performance of broiler chickens. J. Adv. Vet. Anim. Res. 2, 304–309.
34. Janczyk, P., Halle, B. and Souffrant, W.B. 2009. Microbial community composition of the crop and ceca contents of laying hens fed diets supplemented with Chlorella vulgaris. Poult. Sci. 88, 2324–2332.
35. Jeong, J.S. and Kim, I.H. 2014. Effect of Bacillus subtilis C-3102 spores as a probiotic feed supplement on growth performance, noxious gas emission, and intestinal microflora in broilers. Poult. Sci. 93, 3097–3103.
36. Kang, H.K., Salim, H.M., Akter, N., Kim, D.W., Kim, J.H., Bang, H.T., Kim, M.J., Na, J.C., Hwangbo, J., Choi, H.C. and Suh, O.S. 2013. Effect of various forms of dietary Chlorella supplementation on growth performance, immune characteristics, and intestinal microflora population of broiler chickens. J. Appl. Poult. Res. 22, 100–108.
37. Karu, P., Selvan, S.T., Gopi, H. and Manobhavan, M. 2018. Effect of macroalgae supplementation on growth performance of Japanese quails. Int. J. Curr. Microbiol. App. Sci. 7, 1039–1041.
38. Kaushik, P. and Chauhan, A. 2008. In vitro antibacterial activity of laboratory grown culture of Spirulina platensis. Ind. J. Microbiol. 48, 348–352.
39. Lee, C., Kim, J.H. and Kil, D.Y. 2022. Comparison of stress biomarkers in laying hens raised under a long-term multiple stress condition. Poult. Sci. 101(6), 101868; doi: 10.1016/j.psj.2022.101868.
40. Mahrose, K.M., Abol-Ela, S., Amin, R.M. and Abou-Kassem, D.E. 2022. Restricted feeding could enhance feed conversion ratio and egg quality of laying Japanese quail kept under different stocking densities. Anim. Biotechnol. 33, 141–149.
41. Mahrose, K.M., Abd El-Hack, M.E. and Amer, S.A. 2019a. Influences of dietary crude protein and stocking density on growth performance and body measurements of ostrich chicks. An. Acad. Bras. Cienc. 91, e20180479.
42. Mahrose, K.M., Alagawany, M., Abd El-Hack, M.E., Mahgoub, S.A. and Attia, F.A.M. 2019b. Influences of stocking density and dietary probiotic supplementation on growing Japanese quail performance. An. Acad. Bras. Cienc. 91, e20180616.
43. Mahrose, K.M. and Michalak, I. 2022. Seaweeds for animal feed, current status, challenges, and opportunities. In Sustainable global resources of seaweeds, vol. 1. Eds., Rao, A.R. and Ravishankar, G.A. Cham, Switzerland: Springer, pp: 357–359.
44. Mahrose, K.M., Michalak, I., Farghly, M., Elokil, A., Zhang, R., Ayaşan, T., Mekawy, A. and Fazlani, S. 2021. Role of clay in detoxification of aflatoxin B1 in growing Japanese quail with reference to gender. Vet. Res. Comm. 45, 363–371.

45. Mariey, Y.A., Samak, H.R., Abou-Khashba, H.A., Sayed, M.A.M. and Abou-Zeid, A.E. 2014. Effect of using *Spirulina platensis* algae as feed additives for poultry diets: 2-productive performance of broiler. Egypt. Poult. Sci. J. 34, 245–58.
46. Marareni, M., Mhlongo, G. and Mnisi, C.M. 2023. The effect of incorporating dietary green seaweed (*Ulva* sp.) on growth performance, blood parameters, and carcass and meat quality characteristics of Jumbo quail. Heliyon 9, e19603.
47. Michalak, I. and Mahrose, K.M. 2020. Seaweeds, intact and processed, as a valuable component of poultry feed. J. Mar. Sci. Eng. 8, 620; doi: 10.3390/jmse8080620.
48. Maoka, T. 2011. Carotenoids in marine animals. Mar. Drugs. 9, 278–293; doi: 10.3390/md9020278.
49. National Research Council (NRC). 1994. Nutrient requirements of poultry, 9th ed. Washington, D.C., USA: National Academy of Science.
50. Ocak, N. and Erener, G. 2005. The effects of restricted feeding and feed form on growth, carcass characteristics and days to first egg of Japanese quail (*Coturnix coturnix japonica*). Asian-Aust. J. Anim. Sci. 18, 1479–1484.
51. Park, J.H., Lee, S.I. and Kim, I.H. 2018. Effect of dietary *Spirulina* (*Arthrospira*) *platensis* on the growth performance, antioxidant enzyme activity, nutrient digestibility, cecal microflora, excreta noxious gas emission, and breast meat quality of broiler chickens. Poult. Sci. 97, 2451–2459.
52. Rajat, B. 2013. Effects of feed restriction on compensatory growth period, nutrient digestibility and economics of Japanese quails (*Coturnix coturnix japonica*). Int. Multidiscip. Res. J. 3, 4–7.
53. SAS Institute. 1998. SAS user's guide: version 9.1. Cary: SAS Institute Inc.
54. Scholtz, N., Halle, I., Flachowsky, G. and Sauerwein, H. 2009. Serum chemistry reference values in adult Japanese quail (*Coturnix coturnix japonica*) including sex-related differences. Poult. Sci. 88, 1186–1190.
55. Shafiei, A., Khavarinezhad, S., Javandel, F., Nosrati, M., Seidavi, A. and Diarra, S.S. 2018. Effects of duration of early feed withdrawal and re-feeding on growth, carcass traits, plasma constituents and intestinal microflora of broiler chickens. J. Appl. Anim. Res. 46, 1358–1362.
56. Shokri, H., Khosravi, A. and Taghavi, M. 2014. Efficacy of *Spirulina platensis* on immune functions in cancer mice with systemic candidiasis. J. Mycol. Res. 1, 7–13.
57. Sieuwerts, S., De Bok, F.A., Mols, E., De Vos, W.M. and van Hylckama Vlieg, J.E.T.A. 2008. Simple and fast method for determining colony forming units. Lett. Appl. Microbiol. 47, 275–278.
58. Snedecor, G.W. and Cochran, W.G. 1982. Statistical methods, 6th ed. Ames, IA, USA: Iowa State University Press.
59. Soomro, R.N., Abd El-Hack, M.E., Shah, S.S., Taha, A.E., Alagawany, M., Swelum, A.A., Hussein, E.O., Ba-Aawdh, H.A., Saadeldin, I., El-Edel, M.A. and Tufarelli, V. 2019. Impact of restricting feed and probiotic supplementation on growth performance, mortality and carcass traits of meat-type quails. Anim. Sci. J. 90, 1388–1395.
60. Subakir, F.N.M., Ishak, N.I., Samah, A.N., Aziz, K.A.A. and Zaharudin, N. 2021. The effects of seaweed-based pellet binders on growth performance, feed efficiency and carcass characteristics in broilers. Anim. Feed Sci. Technol. 272, 114786; doi: 10.1016/j.anifeedsci.2020.114786.
61. Thiam, M., Wang, Q., Sánchez, A.B., Zhang, J., Ding, J., Wang, H., Zhang, Q., Zhang, N., Wang, J., Li, Q., Wen, J. and Zhao, G. 2022. Heterophil/Lymphocyte ratio level modulates *Salmonella* resistance, cecal microbiota composition and functional capacity in infected chicken. Front. Immunol. 13, 816689; doi: 10.3389/fimmu.2022.816689.
62. Tufarelli, V., Baghban-Kanani, P., Azimi-Youvalari, S., Hosseintabar-Ghasemabad, B., Slozhenkina, M., Gorlov, I., Seidavi, A., Ayaşan, T. and Laudadio, V. 2021. Effects of Horsetail (*Equisetum arvense*) and spirulina (*Spirulina platensis*) dietary supplementation on laying hens productivity and oxidative status. Animals, 11, 335; doi: 10.3390/ani11020335.
63. Tůmová, E., Chodová, D., Härtlová, H., Fučíková, A. and Ketta, M. 2019. Effect of feeding regime on the performance and blood parameters of male and female broiler chickens. S. Afr. J. Anim. Sci. 49, 244–252.
64. Tyl, J., Tůmová, E. and Chodova, D. 2024. The effect of feed restriction and housing system on performance, organ proportion and microbiota. Czech J. Anim. Sci. 69, 68–74.
65. Vijayalingam, T.A., Rajesh, N.V., Vairamuthu, S., Boopathy Raja, M. and Sudeep Kumar, N.K. 2023. Effect of dietary supplementation of seaweeds on growth and blood profiles of TANUVAS Aseel chicken. J. Hellenic Vet. Med. Soc. 74, 6085–6092; doi: 10.12681/jhvms.30812.

66. Wang, S., Hui, J.Y., Hua, W.L., Hua, Z.F. and Ting, L.Y. 2013. Enteromorphaprolifera supplemental level: effects on laying performance, egg quality, immune function and microflora in feces of laying hens. Chinese J. Anim. Nutr. 25, 1346–1352..
<https://doi.org/10.1016/j.indcrop.2019.111992>